

Cycle–Pancyclism in Tournaments II

Hortensia Galeana–Sánchez and Sergio Rajsbaum*

Instituto de Matemáticas

U.N.A.M.

C.U., Circuito Exterior, D.F. 04510, México

(APRIL 17, 1994)

Abstract

Let T be a Hamiltonian tournament with n vertices and γ a Hamiltonian cycle of T . In this paper we develop a general method to find cycles of length k , $\frac{n+4}{2} < k < n$, intersecting γ in a large number of arcs. In particular we can show that if there does not exist a cycle C_k intersecting γ in at least $k - 3$ arcs then for any arc e of γ there exists a cycle C_k containing e and intersecting γ in at least $k - \frac{2(n-3)}{n-k+3} - 2$ arcs. In a previous paper [3] the case of cycles of length k , $k \leq \frac{n+4}{2}$ was studied.

1 Introduction

The subject of pancyclism in tournaments has been studied by several authors (e.g. [1],[2]). Two types of pancyclism have been considered. A tournament T is *vertex-pancyclic* if given any vertex v there are cycles of every length containing v . Similarly, a tournament T is *arc-pancyclic* if given any arc e there are cycles of every length containing e . It is well known that a Hamiltonian tournament is vertex-pancyclic, but not necessarily arc-pancyclic. In a previous paper [3] we introduced the concept of *cycle-pancyclicism* to study questions such as the following. Given a cycle C , what is the maximum number of arcs which a cycle of length k , with vertex set contained in C , has in common with C ? Clearly, to study this kind of questions it is sufficient to consider a Hamiltonian tournament where C is a Hamiltonian cycle of T .

Let T be a Hamiltonian tournament with vertex set $V = \{0, 1, \dots, n - 1\}$ and arc set A . Assume without loss of generality that $\gamma = (0, 1, \dots, n - 1, 0)$ is a Hamiltonian cycle of T . Let C_k denote a directed cycle of length k . In [3] we proved that for $k = 4, 5$ and for every k , such that $n > 2k - 5$, there exists a cycle C_k intersecting γ in at least $k - 3$ arcs. For $k = 3$ it was proved that there exists a cycle C_3 intersecting γ in at least one arc.

*On leave at MIT Laboratory for Computer Science, 545 Technology Square, Cambridge, MA 02139, USA.

In this paper we assume that $k + 1 \leq n \leq 2k - 5$, $k > 5$, and develop methods for finding a cycle C_k intersecting γ in a large number of arcs. In particular we can show that if there does not exist a cycle C_k intersecting γ in at least $k - 3$ arcs then for any arc e of γ there exists a cycle C_k containing e and intersecting γ in at least $k - \frac{2(n-3)}{n-k+3} - 2$. The methods developed in this paper are the basis for our subsequent work in which we study the maximum intersection of a cycle C_k with γ .

2 Preliminaries

A *chord* of a cycle C is an arc not in C with both terminal vertices in C . The *length* of a chord $f = (u, v)$ of C , denoted $l(f)$, is equal to the length of $\langle u, C, v \rangle$, where $\langle u, C, v \rangle$ denotes the uv -directed path contained in C . We say that f is a c -chord if $l(f) = c$ and $f = (u, v)$ is a $-c$ -chord if $l\langle v, C, u \rangle = c$. Observe that if f is a c -chord then it is also a $-(n-c)$ -chord. Unless otherwise stated if the cycle is not specified, it will be assumed that the chord is of cycle γ .

In what follows T is a tournament of n vertices with a Hamiltonian cycle γ . For a cycle C_k of length k with vertex set contained in γ we denote $\mathcal{I}_\gamma(C_k) = |A(\gamma) \cap A(C_k)|$, or simply $\mathcal{I}(C_k)$ when γ is understood. Let $f(n, k, T) = \max\{\mathcal{I}(C_k) \mid C_k \subset T\}$.

Lemma 2.1 *At least one of the following properties holds.*

- (i) $f(n, k, T) \geq k - 3$.
- (ii) *All the following chords are in A .*
 - (a) *Every $(k - 1)$ -chord.*
 - (b) *Every $(k - 2)$ -chord.*

Proof: Suppose that (i) is not true.

(a) If $(k-1, 0)$ is a $-(k-1)$ -chord then $C_k = (0, 1, \dots, k-1, 0)$ satisfies $\mathcal{I}(C_k) = k-1 > k-3$, and hence $f(n, k, T) \geq k-3$.

(b) Suppose that there exists a $-(k-2)$ -chord, $f = (y, x)$. We can assume w.l.o.g. that $x = 1, y = k-1$. It follows from (a) that $(0, k-1) \in A$. Also (a) implies that $(n-(k-1), 0) \in A$ (notation modulo n). Note that the hypothesis $n \leq 2k-5$ implies that $1 < n-(k-1) < k-1$. Let $z \in V$ be the maximum in $\langle 2, \gamma, k-2 \rangle$ such that $(z, 0) \in A$. Since $(0, k-1) \in A$ then $(z, 0) \in A$ and $(0, z+1) \in A$. For $C_k = (z, 0, z+1) \cup \langle z+1, \gamma, k-1 \rangle \cup (k-1, 1) \cup \langle 1, \gamma, z \rangle$ it holds $\mathcal{I}(C_k) = k-3$. ■

3 Lower Bounds

Lemma 3.1 *Let $P = (0, 1, \dots, l)$, $l \geq 2$, be a directed path contained in γ , $z \in V - V(P)$, and $\{(0, z), (1, z), (z, l), (z, l-1), \dots, (z, l-a+1)\} \subseteq A$ with $1 \leq a \leq l-1$. Then there exists i , $0 \leq i \leq l-(a+1)$, such that $\{(i, z), (z, i+a+1)\} \subseteq A$.*

Proof: First we will prove that there exists $j > 1$, such that $j \equiv b \pmod{a+1}$, $b \in \{0, 1\}$, and $(z, j) \in A$. Since $l, l-1, \dots, l-(a-1)$ are consecutive numbers, it follows that there exists $j \in \{l, l-1, \dots, l-(a-1)\}$ with $j \equiv b \pmod{a+1}$ and $b \in \{0, 1\}$; the hypothesis of the lemma implies $(z, j) \in A$, and $j > 1$ because $\{(0, z), (1, z)\} \subseteq A$. Now, let $j_0 = \min\{j | j \equiv b \pmod{a+1}, j > 1, \text{ and } (z, j) \in A\}$. It follows that $(z, j_0 - (a+1)) \notin A$. Hence $(z, j_0) \in A$ and $(j_0 - (a+1), z) \in A$. Clearly, taking $i = j_0 - (a+1)$ we have $\{(i, z), (z, i+a+1)\} \subseteq A$. ■

Lemma 3.2 *If all the $k-2, k-1, \dots, p$ -chords, $k-1 \leq p < n-2$, are in T then at least one of the two following properties hold.*

(i) $f(n, k, T) \geq k-3$.

(ii) Every $(p+1)$ -chord is in T .

Proof: We show that if (ii) is false then (i) holds. Let (s_1, s_2) be a $-(p+1)$ -chord and z a vertex in $\langle s_1, \gamma, s_2 \rangle$. Assume w.l.o.g. that $s_2 = 0$. Let $x \equiv z + n - p \pmod{n}$. Observe that

$$\{(x, z), (x+1, z), \dots, (x+p-(k-2), z)\} \subseteq A \quad (1)$$

since these are the $p, p-1, \dots, k-2$ -chords of γ ending in z . Similarly

$$\{(z, z+p), (z, z+p-1), \dots, (z, z+k-2)\} \subseteq A. \quad (2)$$

Observe that the start points of the arcs in the set (1) are consecutive in γ and less than the end points of the arcs in the set (2), which are also consecutive in γ . This is because the largest start point of an arc in (1) is $z+n-(k-2)$ and the least end point of an arc in (2) is $z+(k-2)$, and $z+(k-2) > z+n-(k-2)$. See Fig. 1.

Now, consider the directed path $\langle x, \gamma, z+p \rangle$. Note that the cardinality of (1) is at least 2 and the cardinality of (2) is $p-k+3$. Thus letting $a = p-k+3$ it follows from Lemma 3.1 that there exist j , $x \leq j < z+(k-2)$ such that $\{(j, z), (z, j+a+1)\} \subseteq A$. It follows that $C = (s_1, s_2) \cup \langle s_2, \gamma, j \rangle \cup (j, z) \cup (z, j+a+1) \cup \langle j+a+1, \gamma, s_1 \rangle$ is a cycle. In order to see that $l(C) = k$ note that $l\langle s_1, \gamma, s_2 \rangle = n - (p+1)$, and thus $l\langle s_2, \gamma, s_1 \rangle = p+1$. Clearly, $l\langle j, \gamma, j+a+1 \rangle = a+1$. Therefore $l(C) = p+1 - (a+1) + 3 = k$ and C is a cycle with $\mathcal{I}(C) = k-3$. ■

It follows directly from Lemma 3.2 the following.

Theorem 3.3 *At least one of the following conditions holds.*

- (i) $f(n, k, T) \geq k - 3$.
- (ii) *For each p , $k - 2 \leq p \leq n - 2$, every p -chord of γ is in T .*

Let l and r be integers such that $n = l(n - k + 3) + r$ and $0 \leq r < n - k + 3$. The following theorem shows that if $f(n, k, T) < k - 3$ then there exist cycles with a large intersection with γ .

Theorem 3.4

- (a) *If $r = 0$ or if $r = 1$ and $k < n - 1$ then $f(n, k, T) \geq k - 2l$.*
- (b) *If $r = 1$ and $k = n - 1$ then $f(n, k, T) \geq k - 2l - 1$.*
- (c) *If $r = 2$ then $f(n, k, T) \geq k - 2l - 1$.*
- (d) *If $r > 2$ then $f(n, k, T) \geq k - 2l - 2$.*

Proof: Notice that for $l = 1$, $0 \leq r \leq 2$ it holds that $3 \leq k \leq 5$. For these cases the theorem follows from [3]. When $l = 1$ we can assume that $r > 2$. Therefore we can assume that the bounds of the theorem $(k - 2l, k - 2l - 1, k - 2l - 2)$ are at most $k - 4$. Therefore if there exists a cycle C_k with $\mathcal{I}(C_k) \geq k - 3$ the theorem follows. Assume that this is not the case, namely, $f(n, k, T) < k - 3$. By Theorem 3.3, for each p , $k - 2 \leq p \leq n - 2$, every p -chord of γ is in T .

We construct a cycle C_k intersecting γ in the required number of arcs. For this goal we specify the following vertices of T , through which C_k passes. Let

$$\begin{aligned} x_1 &= k - 3 \\ x_2 &= x_1 - (n - k + 3) \\ &\vdots \\ x_l &= x_{l-1} - (n - k + 3). \end{aligned}$$

Observe that $l\langle x_{i+1}, \gamma, x_i \rangle = n - k + 3$, and hence by the definition of l , $x_l \geq 0$. It follows from Theorem 3.3 that for every i , $1 \leq i \leq l - 1$, the $(k - 2)$ -chord $(x_i, x_{i+1} + 1)$ is in T . And also, if $z \in V(\gamma)$ such that $2 \leq l\langle x_i, \gamma, z \rangle \leq n - k + 2$, then $(z, x_i) \in T$. For any fixed election of $y_i \in \gamma$, $1 \leq i \leq l - 1$, such that $2 \leq l\langle x_i, \gamma, y_i \rangle \leq n - k + 2$, consider the following path $T_{k'}$ of length k' from 0 to x_{l-1} (Fig. 2)

$$T_{k'} = (0, x_1 + 1) \cup \langle x_1 + 1, \gamma, y_1 \rangle \cup (y_1, x_1, x_2 + 1) \cup \langle x_2 + 1, \gamma, y_2 \rangle \cup (y_2, x_2, x_3 + 1) \cup \cdots \cup (y_{l-1}, x_{l-1}).$$

We now describe how to complete $T_{k'}$ into a cycle C for the different possible values of r . Observe that $l(C)$ will vary depending on the election of the y_i 's, i.e. depending on $l\langle x_i, \gamma, y_i \rangle$.

In each case we prove that C can be constructed in a way such that $l(C) = k$ by a suitable election of y_i , $1 \leq i \leq l-1$, and that $\mathcal{I}(C)$ takes the required values. We shall use the fact that $l(T'_k) \geq 3(l-1)$ and that the number of arcs of T'_k not in γ is $2(l-1)$. Recall that we are assuming that $k > 5$.

- When $r = 0$ ($x_i = 0$) and when $r = 1$ ($x_i = 1$) and $k \leq n-2$, (see Figures 3 and 4)

$$C = T'_k \cup (x_{l-1}, x_l + 1) \cup \langle x_l + 1, \gamma, y_l \rangle \cup (y_l, 0).$$

The reason for not including the case of $k = n-1$ when $r = 1$ is the following. In the description of C the arc $(y_l, 0)$ is assumed to be in T . Theorem 3.3 guarantees the existence of this arc when $y_l \leq x_{l-1} - 2$; when $y_l = x_{l-1} - 1$ the arc $(y_l, 0)$ is a $(k-3)$ -chord. This implies that the vertices $x_{l-1} - 1$ and 1 are not in C and thus k is at most $n-2$.

We proceed to prove that when $r = 0$ and when $r = 1$ ($x_l = 1$) and $k \leq n-2$, $\mathcal{I}(C) = l(C) - 2l$ and C can be constructed to have $l(C) = k$.

The description of C implies that exactly $2l$ arcs of C are not in γ . Therefore $\mathcal{I}(C) = l(C) - 2l$.

Next we show that C can be constructed to have $l(C) = k$. Notice that $C \cap \gamma$ is the union of the paths $\langle x_i + 1, \gamma, y_i \rangle$, for $1 \leq i \leq l$, and hence $l \leq \sum_{i=1}^l l(\langle x_i + 1, \gamma, y_i \rangle) = l(C) - 2l$. Hence, $3l \leq l(C)$. It follows that we can construct cycles C with any $l(C)$, $3l \leq l(C)$, and, by the definition of C , with $l(C) \leq n-1$ ($r = 0$) or $l(C) \leq n-2$ ($r = 1$). Since we want $l(C) = k$ it remains to show that it holds that $3l \leq k$. The proof is as follows. First observe that

$$\frac{3n}{n-k+3} \leq k,$$

because it is equivalent to

$$\begin{aligned} 3n &\leq k(n-k+3) = kn - k^2 + 3k \\ k^2 - 3k - nk + 3n &\leq 0 \\ k(k-3) &\leq n(k-3), \end{aligned}$$

and because $5 < k < n$. Now, if $r = 0$ then $n = l(n-k+3)$. Hence $3l = \frac{3n}{n-k+3}$ and then $3l \leq k$. If $r = 1$ then $n = l(n-k+3) + 1$. Hence $3l = \frac{3(n-1)}{n-k+3} < \frac{3n}{n-k+3} \leq k$.

- When $r = 1$ ($x_i = 1$) and $k = n-1$.

Notice that in this case $y_l = 4$ because $l(\langle x_l, \gamma, x_{l-1} \rangle) = 4$ and $x_{l-1} = 5$, we can define

$$C = T'_k \cup (5, 3, 1, 2, 0),$$

where $y_i = x_{i-1} - 1$ for $1 \leq i \leq l-1$. Clearly $l(C) = n-1$ because the only vertex not in C is 4. The description of C implies that exactly $2l+1$ arcs of C are not in γ . Therefore $\mathcal{I}(C) = k - 2l - 1$.

- When $r = 2$ ($x_l = 2$, Fig. 5)

$$C = T_{k'} \cup (x_{l-1}, x_l + 1) \cup \langle x_l + 1, \gamma, y_l \rangle \cup (y_l, x_l, 0).$$

There are exactly $2l + 1$ arcs of C not in γ . Hence $\mathcal{I}(C) = l(C) - 2l - 1$.

In this case $l(C) \geq 3l + 1$. Since $n = l(n - k + 3) + 2$, to prove that C can be constructed to have $l(C) = k$ it is sufficient to prove that $\frac{3(n-2)}{n-k+3} + 1 \leq k$. The proof is as follows.

$$\begin{aligned} \frac{3(n-2)}{n-k+3} &\leq k-1 \\ 3(n-2) &\leq (k-1)(n-k+3) \\ k^2 - 4k - 3 &\leq n(k-4) \\ k(k-4) &\leq n(k-4), \end{aligned}$$

which holds because $5 < k < n$.

- When $r > 2$ ($x_l > 2$)

$$C = T_{k'} \cup (x_{l-1}, x_l + 1) \cup \langle x_l + 1, \gamma, y_l \rangle \cup (y_l, x_l, 1) \cup \langle 1, \gamma, y_{l+1} \rangle \cup (y_{l+1}, 0).$$

There are exactly $2l + 2$ arcs of C not in γ . Hence $\mathcal{I}(C) = l(C) - 2l - 2$.

In this case $l(C) \geq 3l + 3$. Since $n = l(n - k + 3) + r$, to prove that C can be constructed to have $l(C) = k$ it is sufficient to prove that $3(\frac{n-r}{n-k+3} + 1) \leq k$. Clearly it suffices to prove the inequality for $r = 3$. The proof is as follows.

$$\begin{aligned} 3\left(\frac{n-3}{n-k+3} + 1\right) &\leq k \\ \frac{3(n-3+n-k+3)}{n-k+3} &\leq k \\ \frac{3(2n-k)}{n-k+3} &\leq k \\ 3(2n-k) &\leq k(n-k+3) \\ k(k-6) &\leq n(k-6), \end{aligned}$$

which holds since $5 < k < n$. ■

Notice that Theorem 3.4 guarantees the existence of a cycle C_k with $\mathcal{I}(C_k) \geq k - 2l - 2 \geq k - \frac{2n}{n-k+3} - 2$, for any $(n+5)/2 \leq k \leq n-1$. Therefore, even in the extreme case of $k = n-1$, $f(n, k, T) \geq n/2 - 3$.

Given an arc a of γ , in each of the constructions of the cycle C in the previous theorem, it is possible to construct C in such a way that it contains a . This is done by letting $a = (x_1 + 1, x_1 + 2)$. Hence we have the following

Corollary 3.5 *If $f(n, k, T) < k - 3$ then for any arc $a \in \gamma$ there exists a cycle C_k , $a \in C_k$, with $\mathcal{I}(C_k) \geq k - \frac{2n}{n-k+3} - 2$.*




Figure 1: *Illustrating Lemma 3.2*

References

- [1] B. Alspach, “Cycles of each length in regular tournaments.” *Canadian Math. Bull.*, **10**, (1967), 283–286.
- [2] J.C. Bermond, C.Thomasen, “Cycles in digraphs – A survey.” *J. Graph Theory*, **5**, (1981), 43, 145–157.
- [3] H. Galeana–Sanchez, S. Rajsbaum, “Cycle Pancyclism in Tournaments I,” Pub. Prel. 266 (technical report), Instituto de Matemáticas, UNAM, Mexico, April 1992. Submitted for publication.




Figure 2: *The path T_k'*




Figure 3: When $r = 0$ ($x_l = 0$)




Figure 4: When $r = 1$ ($x_l = 1$) and $k \leq n - 2$




Figure 5: When $r = 2$